

ReproPig®

ReproPig® is used to treat reproductive problems that ultimately enhance the reproductive performance of your breeding herd, both in gilts and sows.



Regumate® Pig

Synchronizes Estrus



EU,
APSA,
LATAM,
NA*



1 Dose
5 ML



Gilts



Sows**

*North America has Altrenogest under the name Matrix, with a different treatment schedule and composition.
**Regumate® is not registered for its use in sows in all the countries.

Gilts

One dose of 5 ml per gilt per day for **18 consecutive days** given orally with feed for immediate consumption.

Sows

Registrations can differ per country. Therefore Regumate may be registered for sows in some countries.

Prepuberal Gilts

P.G. 600 is indicated for induction of fertile estrus (heat) in healthy prepuberal (non-cycling) gilts over **six months of age** and **weighing at least 85 kg (187 lb.)**.

Sows at Weaning

P.G. 600 is indicated for induction of estrus in healthy weaned sows experiencing delayed return to estrus.



PG 600®

Induces Estrus



EU,
APSA,
LATAM,
NA



1 Dose
5 ML



Gilts



Sows



Planate®

Induces Farrowing



1 Dose
2 ML



Gilts



Sows



EU,
APSA,
LATAM,
NA*

*Estrumate (Equine product)

Sow and gilts may not be injected earlier than two days before the expected date to induce the farrowing.

Gilts

Administer 2.5 ml of the product at **115 - 120 hours** after the end of the synchronisation treatment with a progestin. A single artificial insemination should be done **30 - 33 hours** after administration of the product.

Sows

Administer 2.5 ml of the product at **83 - 89 hours** after weaning. A single artificial insemination should be done **30 - 33 hours** after administration of the product.



Porceptal®

Induces Ovulation



1 Dose
2,5 ML



Gilts



Sows



EU,
APSA,
LATAM,
NA