



# SowCare®

The essential range of vaccines for gilts and sows

A complete range of vaccines for the breeding herd, aimed at protecting both sows and their offspring from generation to generation.

## WHEN TO USE

### Porcilis® Ery+Parvo+Lepto

Inactivated vaccine against Porcine Parvovirus, Erysipelas disease and Leptospira.

- EU, APSA, LATAM
- GILTS, SOWS
- 2ml dose volume

**Duration of immunity:**  
*E. rhusiopathiae*: 6 months  
Porcine Parvovirus: 12 months  
*Leptospira* serogroup *Australis*: 6 months  
*Leptospira* serogroups *Canicola*,  
*Icterohaemorrhagiae*, *Grippotyphosa*,  
*Pomona* and *Tarassovi*: 12 months

**Primary vaccination:**  
6 to 8 weeks before insemination

**Booster vaccination:**  
A single vaccination once a year. For *E. rhusiopathiae* containing products and *L. interrogans* serotype *Australis*, revaccination is recommended every 6 months to maintain immunity.

### Porcilis® ColiClos

Inactivated vaccine to provide passive immunity of piglets against *E. coli* and *C. perfringens* type C.

- EU, APSA, LATAM
- GILTS, SOWS
- 2ml dose volume

**Primary vaccination:**  
6 to 8 weeks before farrowing

**Secondary vaccination:**  
4 weeks later

**Booster vaccination:**  
Single vaccination 2 to 4 weeks before farrowing

### Porcilis® Glässer

Inactivated vaccine against *Glaesserella parasuis* infections.

- EU, APSA, LATAM
- SOWS, PIGLETS
- 2ml dose volume

**PIGS**  
**Onset of immunity:**  
2 weeks after completion of vaccination  
**Duration of immunity:**  
14 weeks after completion of vaccination

**SOWS**  
**Onset of immunity:**  
After birth and sufficient uptake of colostrum  
**Duration of immunity:**  
Demonstrated at 4 weeks of age against serotype 4 and at 6 weeks of age against serotype 5

**Vaccination scheme sows and gilts:**  
**Primary vaccination:** Vaccinate sows with the first dose 6 to 8 weeks before farrowing and repeat with a second dose 4 weeks later.  
**Booster vaccination:** Revaccinate 2 to 4 weeks before every subsequent farrowing.

**Vaccination scheme piglets:**  
1st dose at 5 weeks of age, 2nd dose at 7 weeks of age.

### Porcilis® Ery+Parvo

Inactivated vaccine against Erysipelas disease and Porcine Parvovirus infection.

- EU, APSA, LATAM
- GILTS, SOWS
- 2ml dose volume

**Gilts** should be vaccinated against *E. rhusiopathiae* and Porcine Parvovirus before first mating. A single injection, not later than 2 weeks before mating is sufficient to protect the following pregnancy from damage due to Porcine Parvovirus.  
For the induction of protection against erysipelas, an initial course of two vaccinations is recommended. This can be achieved with the single component erysipelas vaccine (eg: Porcilis Ery) given either 4 weeks before, or 4 weeks after the administration of Porcilis Ery+Parvo.

**Sows** should be revaccinated annually with Porcilis Ery + Parvo and this should be supplemented six months later with a single dose of Porcilis Ery vaccine.

### ProSystem® RCE

Modified Live Virus vaccine for the vaccination of healthy pregnant sows, to provide passive protection to their nursing piglets against disease caused by two major Rotavirus serotypes, *E. coli* strains and *C. perfringens* type C.

- APSA, NA
- GILTS, SOWS
- 2ml dose volume

**Primary vaccination:**  
2ml dose IM to pregnant sows and gilts 5 weeks before farrowing

**Secondary vaccination:**  
2ml dose 2 weeks before farrowing

**Booster vaccination:**  
In subsequent farrowings, inject 2ml vaccination dose 2 weeks before farrowing

In subsequent farrowings, inject 1 x 2ml dose 2 to 4 weeks before farrowing

### Porcilis® Porcoli DF

Inactivated vaccine to provide passive immunity for piglets against *E. coli* related diarrhoea and neonatal enterotoxigenesis in first days of life.

- EU, APSA
- GILTS, SOWS
- 2ml dose volume

**Primary vaccination:**  
6 to 8 weeks before farrowing

**Secondary vaccination:**  
4 weeks later

**Booster vaccination:**  
single injection preferably 2 to 4 weeks before next farrowing

In subsequent farrowings, inject 1 x 2ml dose 2 to 4 weeks before farrowing

### Porcilis® Ery

Inactivated vaccine for the control of Erysipelas disease in piglets.

- EU, APSA, LATAM
- GILTS, SOWS
- 2ml dose volume

**Onset of immunity:**  
3 weeks

**Duration of immunity:**  
6 months

**Primary vaccination:**  
a single 2ml dose from 10 weeks of age

**Secondary vaccination:**  
2ml dose 4 weeks later

**Booster vaccination:**  
2ml dose revaccination at 6 monthly intervals

### Porcilis® Parvo

Inactivated vaccine against Porcine Parvovirus related reproductive failure.

- EU, APSA, LATAM
- GILTS, SOWS
- 2ml dose volume

**Onset of immunity:**  
3 weeks

**Duration of immunity:**  
One reproductive cycle

**Gilts:**  
One 2ml dose, 2 weeks before first insemination.

**Sows:**  
One 2ml dose, at least 2 weeks before insemination.

**Booster vaccination:**  
One 2ml dose per year at least 2 weeks before insemination.

### Porcilis® Strepsuis

Active immunisation of pigs against clinical signs due to *Streptococcus suis* serotype 2 infection.

- EU, APSA, LATAM
- PIGLETS
- 2ml dose volume

**Onset of immunity:**  
1 week after the booster vaccination

**Duration of immunity:**  
At least 2 weeks

**Vaccination scheme:**  
Two vaccinations each of 2ml, three weeks apart, from 2 weeks of age.